

Package ‘BenfordTests’

June 5, 2013

Type Package

Title Statistical Tests for Evaluating Conformity to Benford’s Law

Version 1.0.1

Date 2013-06-05

Depends R (>= 3.0.0)

Author Dieter William Joenssen, with contributions from Thomas Muellerleile

Maintainer Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

Description This package contains seven specialized statistical tests and support functions for determining if numerical data could conform to Benford’s law.

License GPL-3

Archs i386, x64

R topics documented:

BenfordTests-package	2
chisq.benftest	3
edist.benftest	5
jpsq.benftest	6
ks.benftest	7
mdist.benftest	9
meandigit.benftest	10
pbenf	11
qbenf	12
rbenf	13
signifd	14
signifd.seq	15
usq.benftest	15
Index	17

BenfordTests-package *Statistical Tests for Benford's Law.*

Description

This package contains seven specialized statistical tests and support functions for determining if numerical data could conform to Benford's law.

Details

Package: BenfordTests
Type: Package
Version: 1.0.1
Date: 2013-06-05
License: GPL-3

BenfordTests is the implementation of the seven most commonly used goodness-of-fit (GOF) tests to assess if data conforms to Benford's law.

Tests include:

Pearson *chi-square* statistic (Pearson (1900))

Kolmogorov-Smirnov *D* statistic (Kolmogorov (1933))

Freedman's modification of Watson's *U-square* statistic (Freedman (1981), Watson (1961))

Chebyshev distance *m* statistic (Leemis (2000))

Euclidean distance *d* statistic (Cho and Gaines (2007))

Judge-Schechter mean deviation *a-star* statistic (Judge and Schechter (2009))

Joenssen's *JP-square* statistic, a Shapiro-Francia type correlation test (Shapiro and Francia (1972))

All tests may be performed using more than one leading digit. All tests simulate the specific p-values required for statistical inference, while p-values for the *chi-square* and *D* statistics may also be determined using their asymptotic distributions.

Author(s)

Dieter William Joenssen

Maintainer: Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Cho WKT, Gaines BJ. Breaking the (Benford) law: Statistical fraud detection in campaign finance. The American Statistician. 2007;61(4):218-223.

Freedman LS. Watson's Un2 statistic for a discrete distribution. Biometrika. 1981;68(3):708-711.

Judge G, Schechter L. Detecting problems in survey data using Benford's law. *Journal of Human Resources*. 2009;44:1-24.

Kolmogorov AN. Sulla determinazione empirica di una legge di distribuzione. *Giornale dell'Istituto Italiano degli Attuari*. 1933;4:83-91.

Leemis LM, Schmeiser BW, Evans DL. Survival distributions satisfying Benford's law. *The American Statistician*. 2000;54(4):236-241.

Newcomb S. Note on the frequency of use of the different digits in natural numbers. *American Journal of Mathematics*. 1881;4(1):39-40.

Pearson K. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. *Philosophical Magazine Series 5*. 1900;50(302):157-175.

Shapiro SS, Francia RS. An approximate analysis of variance test for normality. *Journal of the American Statistical Association*. 1972;67:215-216.

Watson GS. Goodness-of-fit tests on a circle. *Biometrika*. 1961;48:109-114.

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Look at sample
X
#Look at the first digits of the sample
signifd(X)

#Perform a Chi-squared Test on the sample's first digits using defaults
chisq.benftest(X)
#p-value = 0.648
```

chisq.benftest

Pearson's Chi-squared Goodness-of-Fit Test for Benford's Law

Description

chisq.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs Pearson's chi-square goodness-of-fit test to assert if the data conforms to Benford's law.

Usage

```
chisq.benftest(x = NULL, digits = 1, pvalmethod = "asymptotic", pvalsims = 10000)
```

Arguments

<code>x</code>	A numeric vector.
<code>digits</code>	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
<code>pvalmethod</code>	Method used for calculating the p-value. Either "asymptotic" or "simulate".
<code>pvalsims</code>	An integer specifying the number of replicates to use if <code>pvalmethod = "simulate"</code> .

Details

A chi-square goodness-of fit test is performed on `signifd(x,digits)` versus `pbenf(digits)`. `x` is a numeric vector of arbitrary length. Values of `x` should be continuous, as dictated by theory, but may also be integers. `digits` should be chosen so that `signifd(x,digits)` is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

<code>statistic</code>	the value of the chi-squared test statistic
<code>p.value</code>	the p-value for the test
<code>method</code>	a character string indicating the type of test performed
<code>data.name</code>	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Pearson K. On the criterion that a given system of deviations from the probable in the case of a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling. Philosophical Magazine Series 5. 1900;50(302):157-175.

See Also

[pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform a Chi-squared Test on the sample's
#first digits using defaults but determine
#the p-value by simulation
chisq.benftest(X,pvalmethod ="simulate")
#p-value = 0.6401
```

edist.benftest

*Euclidean Distance Test for Benford's Law***Description**

edist.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs a goodness-of-fit test based on the Euclidean distance between the first digits' distribution and Benford's distribution to assert if the data conforms to Benford's law.

Usage

```
edist.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

x	A numeric vector.
digits	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
pvalmethod	Method used for calculating the p-value. Currently only "simulate" is available.
pvalsims	An integer specifying the number of replicates used if pvalmethod = "simulate".

Details

A statistical test is performed utilizing the Euclidean distance between signifd(x,digits) and pbenf(digits). x is a numeric vector of arbitrary length. Values of x should be continuous, as dictated by theory, but may also be integers. digits should be chosen so that signifd(x,digits) is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

statistic	the value of the Euclidean distance test statistic
p.value	the p-value for the test
method	a character string indicating the type of test performed
data.name	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Cho WKT, Gaines BJ. Breaking the (Benford) law: Statistical fraud detection in campaign finance. The American Statistician. 2007;61(4):218-223.

Morrow J. Benford's law, families of distributions and a test basis. 2010.
<http://www.johnmorrow.info/projects/benford/benfordMain.pdf>.

See Also[pbenf](#)**Examples**

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform a Euclidean Distance Test on the
#sample's first digits using defaults
edist.benftest(X,pvalmethod ="simulate")
#p-value = 0.6085
```

jpsq.benftest

*Joenssen's JP-square Test for Benford's Law***Description**

jpsq.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs a goodness-of-fit test based on the correlation between the first digits' distribution and Benford's distribution to assert if the data conforms to Benford's law.

Usage

```
jpsq.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

x	A numeric vector.
digits	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
pvalmethod	Method used for calculating the p-value. Currently only "simulate" is available.
pvalsims	An integer specifying the number of replicates used if pvalmethod = "simulate".

Details

A statistical test is performed utilizing the sign-preserved squared correlation between `signifd(x,digits)` and `pbenf(digits)`. `x` is a numeric vector of arbitrary length. Values of `x` should be continuous, as dictated by theory, but may also be integers. `digits` should be chosen so that `signifd(x,digits)` is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

statistic	the value of the <i>JP-square</i> test statistic
p.value	the p-value for the test
method	a character string indicating the type of test performed
data.name	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Joenssen DW. A new test for Benford's distribution [abstract]. In: Abstract-proceedings of the 3rd joint Statistical Meeting DAGStat, March 18-22, 2013; Freiburg, Germany.

Shapiro SS, Francia RS. An approximate analysis of variance test for normality. Journal of the American Statistical Association. 1972;67:215-216.

See Also

[pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform Joenssen's \emph{JP-square} Test
#on the sample's first digits using defaults
jpsq.benftest(X)
#p-value = 0.3241
```

ks.benftest

Kolmogorov-Smirnov Test for Benford's Law

Description

ks.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs the Kolmogorov-Smirnov goodness-of-fit test to assert if the data conforms to Benford's law.

Usage

```
ks.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

x	A numeric vector.
digits	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
pvalmethod	Method used for calculating the p-value. Currently only "simulate" is available.
pvalsims	An integer specifying the number of replicates used if pvalmethod = "simulate".

Details

A Kolmogorov-Smirnov test is performed between `signifd(x,digits)` and `pbenf(digits)`. `x` is a numeric vector of arbitrary length. Values of `x` should be continuous, as dictated by theory, but may also be integers. `digits` should be chosen so that `signifd(x,digits)` is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

<code>statistic</code>	the value of the Kolmogorov-Smirnov D test statistic
<code>p.value</code>	the p-value for the test
<code>method</code>	a character string indicating the type of test performed
<code>data.name</code>	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Kolmogorov AN. Sulla determinazione empirica di una legge di distribuzione. Giornale dell'Istituto Italiano degli Attuari. 1933;4:83-91.

See Also

[pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform a Kolmogorov-Smirnov Test on the
#sample's first digits using defaults
ks.benftest(X)
#0.7483
```

mdist.benftest	<i>Chebyshev Distance Test (maximum norm) for Benford's Law</i>
----------------	-----------------------------------------------------------------

Description

mdist.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs a goodness-of-fit test based on the Chebyshev distance between the first digits' distribution and Benford's distribution to assert if the data conforms to Benford's law.

Usage

```
mdist.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

x	A numeric vector.
digits	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
pvalmethod	Method used for calculating the p-value. Currently only "simulate" is available.
pvalsims	An integer specifying the number of replicates used if pvalmethod = "simulate".

Details

A statistical test is performed utilizing the Chebyshev distance between `signifd(x,digits)` and `pbenf(digits)`. `x` is a numeric vector of arbitrary length. Values of `x` should be continuous, as dictated by theory, but may also be integers. `digits` should be chosen so that `signifd(x,digits)` is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

statistic	the value of the Chebyshev distance (maximum norm) test statistic
p.value	the p-value for the test
method	a character string indicating the type of test performed
data.name	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Leemis LM, Schmeiser BW, Evans DL. Survival distributions satisfying Benford's law. The American Statistician. 2000;54(4):236-241.

Morrow J. Benford's law, families of distributions and a test basis. 2010.
<http://www.johnmorrow.info/projects/benford/benfordMain.pdf>.

See Also[pbenf](#)**Examples**

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform a Chebyshev Distance Test on the
#sample's first digits using defaults
mdist.benftest(X)
#p-value = 0.6421
```

meandigit.benftest	<i>Judge-Schechter Mean Deviation Test for Benford's Law</i>
--------------------	--------------------------------------------------------------

Description

meandigit.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs a goodness-of-fit test based on the deviation in means of the first digits' distribution and Benford's distribution to assert if the data conforms to Benford's law.

Usage

```
meandigit.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

x	A numeric vector.
digits	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
pvalmethod	Method used for calculating the p-value. Currently only "simulate" is available.
pvalsims	An integer specifying the number of replicates used if pvalmethod = "simulate".

Details

A statistical test is performed utilizing the deviation between the mean digit of signifd(x,digits) and pbenf(digits). The resulting statistic is normalized to [0,1]. x is a numeric vector of arbitrary length. Values of x should be continuous, as dictated by theory, but may also be integers. digits should be chosen so that signifd(x,digits) is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

statistic	the value of the <i>a-star</i> test statistic
p.value	the p-value for the test
method	a character string indicating the type of test performed
data.name	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Judge G, Schechter L. Detecting problems in survey data using Benford's law. Journal of Human Resources. 2009;44:1-24.

See Also

[pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform a Judge-Schechter Mean Deviation Test
#on the sample's first digits using defaults
meandigit.benftest(X)
#p-value = 0.1458
```

pbenf

Distribution Function for Benford's Distribution

Description

Returns the complete Benford distribution function for a given number of first digits.

Usage

```
pbenf(digits = 1)
```

Arguments

digits	An integer determining the number of first digits for which the pdf is returned, i.e. 1 for 1:9, 2 for 10:99 etc.
--------	-------------------------------------------------------------------------------------------------------------------

Value

Returns an object of class "table" containing the expected density of Benford's distribution for the given number of digits.

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

See Also

[qbenf](#); [rbenf](#)

Examples

```
#show Benford's predictions for the frequencies of the first digit values
pbenf(1)
```

qbenf	<i>Quantile Function for Benford's Distribution</i>
-------	-----------------------------------------------------

Description

Returns the complete quantile function for Benford's distribution with a given number of first digits.

Usage

```
qbenf(digits = 1)
```

Arguments

digits	An integer determining the number of first digits for which the qdf is returned, i.e. 1 for 1:9, 2 for 10:99 etc.
--------	-------------------------------------------------------------------------------------------------------------------

Value

Returns an object of class "table" containing the expected quantile function of Benford's distribution with a given number of digits.

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

See Also

[pbenf](#); [rbenf](#)

Examples

```
qbenf(1)

qbenf(1)==cumsum(pbenf(1))
```

rbenf	<i>Random Sample Satisfying Benford's Law</i>
-------	-----------------------------------------------

Description

Returns a random sample with length n satisfying Benford's law.

Usage

```
rbenf(n)
```

Arguments

n	Number of observations.
---	-------------------------

Value

Returns a random sample with length n satisfying Benford's law.

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

See Also

[qbenf](#); [pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Look at sample
X
#should be
# [1] 6.159420 1.396476 5.193371 2.064033 7.001284 5.006184
#7.950332 4.822725 3.386809 1.619609 2.080063 2.242473 1.944697 5.460581
#[15] 6.443031 2.662821 2.079283 3.703353 1.364175 3.354136
```

`signifd`*Leading Digits*

Description

Returns the specified number of significant digits for each element of a given vector.

Usage

```
signifd(x = NULL, digits = 1)
```

Arguments

<code>x</code>	A numeric vector.
<code>digits</code>	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.

Details

`x` is a numeric vector of arbitrary length. Unlike other solutions, this function will work reliably with all real numbers.

Value

Returns a vector of integers the same length as the input vector `x`.

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

See Also

[chisq.benftest](#); [ks.benftest](#); [usq.benftest](#); [mdist.benftest](#); [edist.benftest](#); [meandigit.benftest](#); [jpsq.benftest](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Look at the first digits of the sample
signifd(X)
#should be:
#[1] 6 1 5 2 7 5 7 4 3 1 2 2 1 5 6 2 2 3 1 3
```

signifd.seq*Sequence of Possible Leading Digits*

Description

Returns a vector containing all possible significant digits for a given number of places.

Usage

```
signifd.seq(digits = 1)
```

Arguments

digits An integer determining the number of first digits to be returned, i.e. 1 for 1:9, 2 for 10:99 etc.

Value

Returns an integer vector.

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

Examples

```
signifd.seq(1)
seq(from=1,to=9)==signifd.seq(1)

signifd.seq(2)
seq(from=10,to=99)==signifd.seq(2)
```

usq.benftest*Freedman-Watson U-squared Test for Benford's Law*

Description

usq.benftest takes any numerical vector reduces the sample to the specified number of significant digits and performs the Freedman-Watson test for discreet distributions between the first digits' distribution and Benford's distribution to assert if the data conforms to Benford's law.

Usage

```
usq.benftest(x = NULL, digits = 1, pvalmethod = "simulate", pvalsims = 10000)
```

Arguments

<code>x</code>	A numeric vector.
<code>digits</code>	An integer determining the number of first digits to use for testing, i.e. 1 for only the first, 2 for the first two etc.
<code>pvalmethod</code>	Method used for calculating the p-value. Currently only "simulate" is available.
<code>pvalsims</code>	An integer specifying the number of replicates used if <code>pvalmethod = "simulate"</code> .

Details

A Freedman-Watson test for discrete distributions is performed between `signifd(x,digits)` and `pbenf(digits)`. `x` is a numeric vector of arbitrary length. Values of `x` should be continuous, as dictated by theory, but may also be integers. `digits` should be chosen so that `signifd(x,digits)` is not influenced by previous rounding.

Value

A list with class "htest" containing the following components:

<code>statistic</code>	the value of the <i>U-square</i> test statistic
<code>p.value</code>	the p-value for the test
<code>method</code>	a character string indicating the type of test performed
<code>data.name</code>	a character string giving the name of the data

Author(s)

Dieter William Joenssen <Dieter.Joenssen@TU-Ilmenau.de>

References

Benford F. The law of anomalous numbers. Proceedings of the American Philosophical Society. 1938;78:551-572.

Freedman LS. Watson's U^2 statistic for a discrete distribution. Biometrika. 1981;68(3):708-711.

Watson GS. Goodness-of-fit tests on a circle. Biometrika. 1961;48:109-114.

See Also

[pbenf](#)

Examples

```
#Set the random seed to an arbitrary number
set.seed(421)
#Create a sample satisfying Benford's law
X<-rbenf(n=20)
#Perform Freedman-Watson U-squared Test on
#the sample's first digits using defaults
usq.benftest(X)
#p-value = 0.4847
```


Index

*Topic **datagen**

BenfordTests-package, [2](#)
rbenf, [13](#)

*Topic **distribution**

BenfordTests-package, [2](#)
pbenf, [11](#)
qbenf, [12](#)
rbenf, [13](#)

*Topic **htest**

BenfordTests-package, [2](#)
chisq.benftest, [3](#)
edist.benftest, [5](#)
jpsq.benftest, [6](#)
ks.benftest, [7](#)
mdist.benftest, [9](#)
meandigit.benftest, [10](#)
usq.benftest, [15](#)

*Topic **manip**

BenfordTests-package, [2](#)
signifd, [14](#)

BenfordTests (BenfordTests-package), [2](#)

BenfordTests-package, [2](#)

chisq.benftest, [3](#), [14](#)

edist.benftest, [5](#), [14](#)

jpsq.benftest, [6](#), [14](#)

ks.benftest, [7](#), [14](#)

mdist.benftest, [9](#), [14](#)

meandigit.benftest, [10](#), [14](#)

pbenf, [4](#), [6–8](#), [10](#), [11](#), [11](#), [12](#), [13](#), [16](#)

qbenf, [12](#), [12](#), [13](#)

rbenf, [12](#), [13](#)

signifd, [14](#)

signifd.seq, [15](#)

usq.benftest, [14](#), [15](#)